



Review

Toscana virus: A comprehensive review of 1381 cases showing an emerging threat in the Mediterranean regions

Nazli Ayhan^{a,b,*}, Carole Eldin^{a,c}, Remi Charrel^{a,c,d}^a Unité des Virus Émergents (UVE: Aix-Marseille Univ, Université di Corsica, IRD 190, Inserm 1207, IRBA), France^b Centre National de Référence des Arbovirus, Inserm-IRBA, Marseille, France^c Laboratoire des Infections Virales Aigües et Tropicales, Pôle des Maladies Infectieuses, AP-HM Hôpitaux Universitaires de Marseille, France^d Le Service de Prévention du Risque Infectieux (LESPRI), CLIN AP-HM Hôpitaux Universitaires de Marseille, France

ARTICLE INFO

Article history:

Accepted 12 January 2025

Available online 17 January 2025

Keywords:

Phlebovirus

Toscana virus

Clinical characterization

Arbovirus

Phlebovirus toscanaense

SUMMARY

Background: Toscana virus (TOSV) is a sand fly-borne phlebovirus causing central nervous system (CNS) infection in Mediterranean countries, during summer season. However, clinical aspects of the disease caused by this virus are poorly known by clinicians, so that its prevalence is probably underestimated due to a lack of diagnosis.

Study design: The data was gathered from all available case series and retrospective studies identifying TOSV as the causative viral agent. The informations of age, sex, clinical characteristics, laboratory findings, imaging results and clinical outcomes of TOSV infection were recorded and analyzed.

Results: A total of 95 articles including TOSV infections resulting in a total of 1381 cases, were analyzed. Our findings indicate that TOSV affects individuals across various age groups, with a median age of 44.45 years. A notable disparity in infection rates between genders, with men being significantly more likely to present symptoms due to TOSV than women, with a sex ratio of 2.0. The clinical presentation of TOSV infection encompasses a range of symptoms, including fever, headache, retro-orbital pain, neurological and muscular manifestations with less common reports of cutaneous and gastrointestinal symptoms. To date, six fatalities have been attributed to TOSV infections, with a median age of 76 years.

Diagnostic evaluation of TOSV infections often involves the analysis of cerebrospinal fluid, where findings may include an elevated white blood cell count.

Conclusions: These findings underscore the diverse clinical manifestations of TOSV infections including flu like symptoms. TOSV is an emerging infectious threat that warrants inclusion in the diagnostic protocols for patients presenting with CNS, particularly within the Mediterranean basin or for those with recent travel history to endemic regions during warmer months when sand flies are actively circulating.

© 2025 The Author(s). Published by Elsevier Ltd on behalf of The British Infection Association. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

Toscana virus (TOSV) is classified within the *Phlebovirus toscanaense* species of the *Phenuiviridae* family within the *Bunyavirales* order.¹ TOSV is recognized as a significant human pathogen prevalent in the Mediterranean region.² TOSV is primarily transmitted by sand flies of the *Phlebotomus* (*Ph.*) genus, particularly *Ph. perniciosus* and *Ph. perfiliewi*.³ The vertical transmission between female *Ph. perniciosus* to her offsprings have been showed previously.⁴ The specific animal reservoir(s) for TOSV have not been identified yet. Recent studies, including Medlock

et al.⁵ have documented a concerning increase in the population density and species number of sand flies, along with their dissemination into territories previously considered free of sand flies like Austria in the last decade. The geographical expansion of the sand fly population substantially increases the risk of TOSV exposure among humans. Cases of TOSV infection have been reported in many Mediterranean countries such as Italy, Spain, Portugal, France, Türkiye, Croatia, Greece, Algeria and Tunisia.⁶ Seroprevalence studies in both human and non-human vertebrates have revealed significant infection rates, with high prevalence reported in Italy (19.8%),⁷ Türkiye (17.8%),⁸ Greece (21%),⁹ North Africa (22–41%)¹⁰; and the Balkans (37.5%)¹¹ indicating the widespread presence of TOSV across the Mediterranean basin. In a recent retrospective study in Germany, Dersch et al. reported cases of TOSV neuroinvasive disease in patients without recent travel to endemic regions, indicating a potentially broader geographical distribution of the virus than previously recognized.¹² However, the rare inclusion of TOSV in the diagnostic

* Correspondence to: Unité des Virus Émergents (UVE) Méditerranée Infection, 19–21, Bd. Jean Moulin, 13005 Marseille, France.

E-mail addresses: nazli.ayhan@univ-amu.fr (N. Ayhan), carole.eldin@univ-amu.fr (C. Eldin), remi.charrel@univ-amu.fr (R. Charrel).

algorithm of nervous system infections (CNS) infections likely results in an important underestimation of the incidence of TOSV cases.⁶

The objective of our study was to offer a descriptive analysis of published data about clinical characteristics of TOSV infections to raise the awareness of physicians about this emerging pathogen.

Material and method

Study design

Relevant entries in global Web-based resources that comprise Scopus (<http://www.scopus.com/>), Web of Science (<https://isiknowledge.com>), and PubMed (<https://pubmed.ncbi.nlm.nih.gov>), Google Scholar (<https://scholar.google.com.tr/>) were searched. Database investigations were performed using the keywords “bunyavirus”, “phlebovirus”, “Toscana virus”, “TOSV”, “Toscana virus case report”, “nervous system infection”. There were not any language restriction; the articles in German, Spanish and Italian were also included the data based. The time frame extended from the first reported TOSV case in 1985 to 2023. Reports unrelated to TOSV infection were omitted, as well as conference reports with recurring data in publications. The classification of TOSV diagnoses and methods and clinical classification of neurological manifestations such as meningitis/encephalitis relied on techniques from published articles. The references cited in each report were examined for further publications, which were included in the analysis (Fig. 1).

Statistical analysis

All variables (described in Table 1) were screened using a logistic regression univariate analysis to assess for statistically significant associations of demographic characters with TOSV in SPSS version 24. Each predictor is analyzed separately to see if it has a significant

Table 1
Demographic characteristics and non-specific clinical signs.

	Number of cases	Total ^a	%
Age, median [range]	44.45	762	
Sex (male/female)	744/371	1,115	
Signs ^b			
Retro-orbital pain	156	163	95.7
Headache	566	607	93.2
Fever	678	738	91.9
Neck rigidity	473	536	88.2
Nausea/Vomiting	340	431	78.9
Muscle weakness / Asthenia / Fatigue	34	43	79.1

^a For demographic characteristics, the total number is determined by collecting the available data.

^b For clinical signs, the total number is calculated based on the reported presence or absence sign in the screened sources.

association with the binary outcome of infection (infected vs. not infected). The missing data is excluded from the analysis.

Results

Demographic and geographic characteristics

We conducted an inventory of 95 articles including 48 case reports, 32 case series and 15 retrospective studies, documenting a total of 1381 cases of TOSV infection cases between 1985 and 2023. Age and sex demographics were obtained for 762 and 1115 patients, respectively (Table 1). More male patients were reported with TOSV infection compared to female patients (744 males to 371 females; $p < 0.001$).

TOSV infections were recorded in 12 countries: with the majority of cases originating from Italy ($n=1064$), followed by Spain ($n=76$), Greece ($n=45$), Türkiye and Tunisia each with $n=31$, France ($n=27$), Algeria ($n=23$), and lesser numbers from Croatia ($n=12$), Portugal




Fig. 1. TOSV cases screening diagram.




Fig. 2. Toscana virus local and imported case distribution map between 1985 and 2023.

(n=12), Romania (n=8), Bosnia & Herzegovina (n=7), Germany (n=4) and Israel (n=1). A total of 40 travelers returning from Mediterranean countries were reported, of which 30 had visited Italy, including both mainland and islands (Fig. 2).

TOSV infections were reported across a wide range of age groups, with the youngest patient being between 0–5 year old¹³ and the oldest between 90–95 year old¹⁴; the median age of patients was 44.4 years. The disease manifested in 82 pediatric patients under 15 years of age, as documented in nine studies.^{13–21} Notably, the prevalence of TOSV infection in children was lower, at 5.9%.

Clinical characteristics

Typical manifestations

Retro-orbital pain was the most common symptom (156/163, 95.7%). The headache was observed in 93.2% of cases (566/607) sometimes reported as the “worst headache of their life”.²² Fever was described in 91.9% (678/738) of cases with an abrupt onset and temperature ranging from 38 °C to 39.5 °C. Neck rigidity was reported in 88.2% (473/536). Other non-specific signs such as nausea/vomiting, muscle weakness, asthenia and fatigue were described in almost 80% of cases (Table 1).

Neurological manifestations

Of the 644 documented cases, 519 patients (representing 80.6%) presented with meningitis or aseptic meningitis, while 111 patients presented with either pure encephalitis or meningo-encephalitis (Table 2); myelitis was very rarely reported. Until recently, a total of

Table 2

Number of TOSV cases with muscular, ocular, gastro-intestinal, cutaneous and neurological manifestations.

	Number of cases	Total ^a	%
Muscular Manifestations	51	211	24.2
Myalgia / Arthralgia	51	211	24.2
Ocular Manifestations	157	164	95.7
Retroorbital pain or pressure	156	163	95.7
Gastro-intestinal Manifestations	16	43	37.2
Gastroenteritis	7	16	43.8
Cutaneous Manifestations	15	38	39.5
Rash	11	32	34.4
Neurological Manifestations	569	712	79.9
Central Nervous System Manifestations	560	683	82.0
Peripheral Nervous System Manifestations	14	26	53.8
Meningitis /Aseptic Meningitis	519	644	80.6
Encephalitis	77	241	32.0
Meningoencephalitis	34	221	15.4
Meningitis/Aseptic Meningitis or Meningoencephalitis	49		
Myelitis	2		
Total of Neurological Manifestations	684+331 ^b = 1.015	1.382	73.44

The article did not provide specific patient counts for each clinical presentation due to a lack of detailed information, those numbers were only included in the overall total.²¹

^a The total number is determined based on the reported presence or absence of symptoms.

^b However, a recent article included a total of 331 patients with meningitis, meningoencephalitis, encephalitis, and polyneuropathy.

Table 3
TOSV infection cases with testicular manifestations.

Characteristic / References	Age range / Age median [range] (n)	Orchitis	Testicular swelling and tenderness	Testicular pain	Fever	Headache	Neck rigidity / pain	Nausea / vomiting
Echevarria et al. 2003	25–29	1	-	-	NR	NR	NR	NR
Baldelli et al. 2004	15–19	1	-	-	1	1	1	1
Zanelli et al. 2013	25–29	-	1	1	1	1	-	-
Tschumi et al. 2019	20–24	-	1	1	1	1	-	-
Mascitti et al. 2020	40–44	-	-	1	1	1	-	-
Total / Median age [range] (n)	27 ^{16–44} (5)	2	2	3	4	4	1	1

NR, not reported.

49 cases were recorded as meningitis/meningo-encephalitis. More recently, Mellace et al.²¹ identified 331 cases featuring neurological symptoms; however, the distinction between meningitis and encephalitis/meningoencephalitis was not clearly defined. As a consequence, although central manifestations are not unusual, pure meningitis is the most prominent neurological manifestation of TOSV infections.

Three cases of hydrocephalus were documented in young patients (range between 15–25 year-old); in all cases, hydrocephalus was observed as a complication following viral meningoencephalitis.^{23,24}

Specific neurological manifestations were categorized based on their involvement of either the central or peripheral nervous system. Signs and symptoms denoting a central nervous system involvement was included abnormal behavior, agitation, anhedonia irritability, aphasia, ataxia, blurred vision, cerebral atrophy, confusion, depression, double vision, dysmetria, epilepsy, facial paresis, focal neurological deficit, hallucinations, hemiparesis/paresis, hydrocephalus, hypotonia, impaired consciousness, insomnia, nystagmus, obtundation, occipital/ischemic stroke, paresis, paresthesia, phonophobia, photophobia, kernig's sign, seizure, sleepiness, stumbling and dysmetria, tetra paresis, tonic-clonic seizure, tremor. And signs and symptoms denoting a peripheral nervous system involvement included brachial plexitis, Guillain Barré like syndrome, hyperaesthesia, limited atrophy, motor and sensory neuropathy, peripheral polyneuropathy.

Out of 712 patients, 569 exhibited at least one of the neurological signs or symptoms. Among central neurological manifestations, Kernig's sign was the most prevalent, observed in 135 cases, followed by decreased consciousness and photophobia (n=41), facial or leg hemiparesis /paresis (n=17) and confusion (n=16). Speech impairment was reported in eight patients.^{25–31} Six patients experienced hearing impairment, characterized by bilateral deafness that persisted beyond the acute phase.^{32–35} Additionally, changes in personality, sexual and social disinhibition, aggressiveness, and other abnormal behaviors were also documented^{36,37} (Table 2). The presence of symptoms signing a peripheral neuropathy was observed in 14 cases of which 10 presented with Guillain-Barre-like syndrome.^{38–40}

Ocular manifestations

Retro-orbital pain or pressure was very frequently observed, documented in 156 out of 163 cases. In contrast, conjunctivitis was reported only once⁴¹ (Table 2).

Gastro-intestinal manifestations

Gastro-intestinal manifestations consisting of gastroenteritis, abdominal pain, dysphagia or diarrhea were reported rarely (16/43).^{23,40,42–44} (Table 2).

Cutaneous manifestations

Dermatological symptoms including petechiae, rash, exanthema, and febrile erythema were reported in 15 out of 38 cases. Rash was the most frequently observed symptom, noted in 11 cases^{23,32,42,44–49} (Table 2).

Muscular manifestations

The musculoskeletal symptoms included cramps, myalgia/arthritis, myositis/fasciitis, and muscle stiffness. Among the 51 patients who exhibited these symptoms, all presented with myalgia and/or arthralgia^{14,20,25,38,41,47,48,50–56} (Table 2).

Testicular manifestations

Testicular manifestations, including epididymo-orchitis, tenderness, testicular pain, and swelling, were reported in five patients^{23,42,44,57,58} (Table 3). These symptoms were significantly more frequent (p-value < 0.001) among younger patients, with a median age of 27 years (range 16–44) (Table 3). Recently, TOSV RNA was detected in the seminal fluid of a patient in the age range of 20–25 year old until day 59 after infection without any testicular manifestations.³¹

Severe cases and mortality

Although TOSV infections are frequently associated with severe clinical manifestations, the vast majority of patients recover completely. Nonetheless, six fatal cases have been reported across two studies: the first fatality occurred in Italy³⁴; and five subsequent fatalities were reported in Romania.^{34,52} The overall case fatality rate is 0.43% with deceased patients having a median age of 76 years-old (range 70–91). Age was found to have a statistically significant impact on mortality (p-value < 0.001). In addition, five out of six fatal cases had comorbidities such as hypertension and diabetes (Table 4).

There were nine cases of severe TOSV infections leading to coma.^{23,34,52,59} Seven of these were elder patients with median age of 79, six of whom had comorbidities such as diabetes and hypertension and two patients were two siblings with the age range between 15 to 20, previously mentioned as having developed hydrocephalus following infection.

Laboratory characteristics

Cerebrospinal fluid (CSF) analysis showed elevated white blood cell (WBC) levels in 22 cases. Lymphocyte ratio was high in 82% of the patients (47 out of 57) (> 50%). CSF samples showed high protein levels in 102 patients. Mildly elevated glucose levels have been showed in 179 patients (Table 5).

Electroencephalogram analysis

The electroencephalogram (EEG) showed abnormal results in 43 patients^{13,26,34,59–65} with non-specific abnormalities, occasional spikes and slow waves.⁶¹

Imaging and radiological findings and electrocardiogram tests

When available, magnetic resonance imaging (MRI) results showed diffuse encephalopathy, diffuse atrophy abnormalities, diffuse bilateral asymmetric myositis or scattered punctuate T2-hyperintense non-enhancing white matter lesions in 11 patients.^{24,30,48,56,66} Cerebral atrophy and/or occipital stroke sequelae were detected in 3 patients with computerized tomography (CT) scan.⁵²

Table 4
TOSV infection fatal cases.

Reference	Sex	Age Range	Country	Comorbidities	Period	In-hospital length of stay (days)
Bartels et al. 2012	male	70–74	Italy	-	-	14
Popescu et al. 2021	male	90–94	Romania	Hypertension, congestive heart failure, ischemic heart disease, stroke sequelae	June	10
	female	65–69	Romania	Hypertension, diabetes mellitus, obesity, NonHodgkin lymphoma	August	4
	male	75–79	Romania	Diabetes mellitus type II, Ischemic heart disease	August	19
	male	70–74	Romania	Diabetes mellitus type II, Ischemic heart disease, Atrial fibrillation, Congestive heart failure, Diabetic polyneuropathy and arteriopathy	August	4
	female	85–89	Romania	Hypertension, Stroke sequelae, Chronic renal failure	August	6

Table 5
TOSV infection laboratory characteristics.

Laboratory Findings	Mean (n) [range]	Normal range/percentage
CSF protein level (mg/dL)	56.3 (77) [11–757]	15–45
CSF WBC (cells/mm ³)	432.5 (22) [6–3500]	0–5
CSF Lymphocyte ratio (%)	73.4% (57) [18%–100%]	50%
CSF Glucose (mg/dL)	63.2 (60) [29–132]	> 40

Few patients were examined by with electrocardiogram (ECG) and chest X-ray. Chest X-ray demonstrated an opacity in the right middle lobe of the lung of one patient.³² No abnormalities were recorded with ECG after TOSV infection.

Hospitalization length and symptom duration

Both length of hospitalization and symptoms duration vary significantly based on the severity of the case. The longest hospitalization was 60 days⁶⁷ and the longest symptoms duration recorded as 28 days.^{55,68}

Discussion

Infections caused by TOSV exhibit a high diversity of clinical manifestations, none of them being neither pathognomonic nor highly specific. Unlike other neuroinvasive viruses such as herpes viruses and enteroviruses, TOSV infections occur exclusively during the period of activity of phlebotomine sand flies, typically from March to November in the Mediterranean, with slight variations depending on local climate conditions.²²

From 1985 to 2023, a total of 1381 cases were reported from 12 different countries of which the majority (1064 cases, 77%) occurred in Italy. The annual incidence rate of cases fluctuates considerably in relation with complex biology of vectors, potential hosts and environmental factors. TOSV infections have been documented across a broad age spectrum (median age 44.45 years, ranging from 0 to 95 years old); however, age itself is not indicative of etiology, though most cases occur between 20 and 60 years old. Based on data collected from returning travelers, the median incubation period for the neuroinvasive forms was calculated to be 12 days (range: 10–14 days) which is quite long.⁶⁹ It remains unclear whether the same incubation period applies to cases with mild signs and symptoms such as only fever or headache.

TOSV symptomatic infections are observed twice as frequently in males as in females. The reasons for this gender disparity in symptomatology are not elucidated. It could be related to differences in immune response, viral susceptibility, or possibly due to variations in exposure factors and the prevalence of high-risk behaviors. However, similar seroprevalence rates between males and females suggest that behavioral factors alone may not explain this difference.⁷⁰ Interestingly, the same tendency is also observed for WNV infections.⁷¹

The case fatality rate of TOSV infections is 0.43% which is lower than other neuroinvasive arboviruses such as WNV, Japanese encephalitis virus (JEV) or tick-borne encephalitis virus (TBEV).^{71–73} All the fatal cases were older than 60 year-old; five of the six fatal cases had comorbidities such as hypertension and/or diabetes mellitus.

Severe neuroinvasive forms, characterized by encephalitis or meningo-encephalitis, are reported in 15.8 to 32.0% of neurological cases, the latter representing almost 80% of 712 studied cases. Since TOSV cases reported in the literature are the most severe ones, neuroinvasive forms and mortality rate are certainly over-represented. Mild cases are frequent, although they are likely to remain either undetected or unpublished⁷⁴; those forms are associated with unspecific clinical manifestations such as fever, headache, nausea, vomiting, retro-orbital pain and muscle weakness.

Severe long-lasting or permanent neurological sequelae such as consciousness impairment, hydrocephalus, ischemic stroke and aphasia^{23,52,67,75,76} were described. In contrast with other arboviruses, severe neurological forms are not more frequent in the elderly.²¹ However, the lower incidence rate of TOSV infections in the pediatric group (5.9%) can either be explained by a lower exposure and/or by a reporting bias, or by biological factors such as immune response and/or viral susceptibility.

Among system-specific manifestations, CNS signs are the most frequently observed with 79.9% of cases followed by ocular and muscular manifestations. Peripheral neurological manifestations, gastro-intestinal and cutaneous manifestations were reported rarely. Additionally, a unique case involved a patient with gastrointestinal manifestations who was also co-infected with West Nile virus.⁵⁰ Although testicular manifestations are noteworthy, they have been documented in only five patients. To date, aside from the Zika virus (ZIKV), no other arbovirus has been identified as a causative agent for testicular manifestations. Recently, the presence and persistence of TOSV RNA in seminal fluid has been demonstrated without evidence for sexual transmission.³¹

In most of cases presenting with clinical picture justifying CSF collection, CSF was clear and colorless. The CSF formula showed elevated WBC levels in 22 patients and lymphocytic meningitis in 58 patients. Mildly elevated protein and glucose levels were described in 102 patients and 179 patients respectively. Hypoglycorrhachia was never reported. CSF parameters were available only for a subset of the reviewed cases.

Electro-physiology analysis and radiology findings showed abnormalities in EEG or in MRI for 56 cases (4%). CT scan and chest X-ray were performed occasionally and abnormalities were rarely recorded. No abnormalities were observed in the ECG, which was conducted for only three patients.^{51,77,78}

In almost all cases, the recovery is complete without persisting functional sequelae. However, there are a few documented cases where individuals have experienced complicated forms of meningitis and/or encephalitis with lingering effects attributed to TOSV. These effects include impaired cognitive functions and altered social and sexual behaviors (60,68) Other neuroinvasive arboviruses, such as WNV, JEV or TBEV, cause long-lasting or permanent sequelae.⁷⁹

Finally, most TOSV infections remain undiagnosed, as TOSV is not included in the list of pathogens to be screened in patients

presenting with febrile illness and/or neurological manifestations in areas where the virus is endemic. The only exception is some regions in Italy (Emilia-Romagna) where it is recommended to include TOSV in the panel of viruses to be tested during summertime for suspect cases.²¹ This is likely why most cases are reported in Italy besides the fact that TOSV has been discovered in Italy and that physicians are much more aware of its existence than in other at-risk countries. This oversight is highlighted by numerous retrospective studies, which suggest that many cases of TOSV infection are classified as "infections due to unknown pathogenic agents" due to the absence of specific laboratory screening for TOSV.^{38,54,80,81} Nevertheless, most TOSV cases exhibit nonspecific signs or symptoms with short duration in non-severe cases.

The lack of physician awareness and the absence of recommendations for screening suspected cases using specific laboratory tests have significantly contributed to the underestimation of TOSV cases, despite its importance to public health.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

This study was funded by European Commission grant 101057690 and UKRI grants 10038150 and 10039289, and is catalogued by the CLIMOS Scientific Committee as CLIMOS number 016 (<http://www.climos-project.eu>). The contents of this publication are the sole responsibility of the authors and do not necessarily reflect the views of the European Commission, the Health, and Digital Executive Agency, or UKRI. Neither the European Union nor granting authority nor UKRI can be held responsible.

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. For the purposes of Open Access, the authors have applied a CC BY [option: CC BY-ND] public copyright license to any Author Accepted Manuscript version arising from this submission.

CLIMOS forms part of the Climate Change and Health Cluster comprising six Horizon Europe projects: BlueAdapt, CATALYSE, CLIMOS, HIGH Horizons, IDAlert, and TRIGGER.

Appendix A. Supporting information

Supplementary data associated with this article can be found in the online version at [doi:10.1016/j.jinf.2025.106415](https://doi.org/10.1016/j.jinf.2025.106415).

References

- Kuhn JH, Abe J, Adkins S, Alkhovsky SV, Avšič-Županc T, Ayllón MA, et al. *Annual (2023) taxonomic update of RNA-directed RNA polymerase-encoding negative-sense RNA viruses (realm Riboviria: kingdom Orthornavirae: phylum Negarnaviricota)*. *J Gen Virol* 2023 Aug; **104**(8):001864.
- Charrel RN, Bichaud L, de Lamballerie X. Emergence of Toscana virus in the mediterranean area. *World J Virol* Oct. 2012; **1**(5):135–41.
- Ayhan N, Prudhomme J, Laroche L, Bañuls AL, Charrel RN. Broader geographical distribution of toscana virus in the mediterranean region suggests the existence of larger varieties of sand fly vectors. *Microorganisms* 2020; **8**(1):114.
- MAROLI M, CIUFOLINI MG, VERANI P. Vertical transmission of Toscana virus in the sand fly, *Phlebotomus perniciosus*, via the second gonotrophic cycle. *Med Vet Entomol* Jul. 1993; **7**(3):283–6.
- Medlock JM, Hansford KM, Van Bortel W, Zeller H, Alten B. A summary of the evidence for the change in European distribution of phlebotomine sand flies (Diptera: Psychodidae) of public health importance. *J Vector Ecol* Jun. 2014; **39**(1):72–7.
- Ayhan N, Charrel RN. An update on Toscana virus distribution, genetics, medical and diagnostic aspects. *Clin Microbiol Infect* 2020; **26**(8):1017–23.
- Terrosi C, Olivieri R, Bianco C, Cellesi C, Cusi MG. Age-dependent seroprevalence of Toscana virus in central Italy and correlation with the clinical profile. *Clin Vaccine Immunol* Aug. 2009; **16**(8):1251–2.
- Ergunay K, Aydoğan S, İlhami Özcebe O, Cilek EE, Hacıoğlu S, Karakaya J, et al. *Toscana Virus (TOSV) exposure is confirmed in blood donors from Central, North and South/Southeast Anatolia, Turkey*. *Zoonoses Public Health* Mar. 2012; **59**(2):148–54.
- Anagnostou V, Papa A. Seroprevalence of Toscana virus among residents of Aegean Sea islands, Greece. *Travel Med Infect Dis* Mar. 2013; **11**(2):98–102.
- Alkan C, Allal-Ikhlef AB, Alwassouf S, Baklouti A, Piorkowski G, de Lamballerie X, et al. Virus isolation, genetic characterization and seroprevalence of Toscana virus in Algeria. *Clin Microbiol Infect* Nov. 2015; **21**(11):1040.e1–9.
- Punda-Polic V, Jerončić A, Mohar B, Kraljević KS. Prevalence of Toscana virus antibodies in residents of Croatia. *Clin Microbiol Infect* 2012; **18**(6):200–3.
- Dersch R, Sophocleous A, Cadar D, Emmerich P, Schmidt-Chanasit J, Rauer S. Toscana virus encephalitis in Southwest Germany: a retrospective study. *BMC Neurol* Dec. 2021; **21**(1):1–5.
- Braito A, Ciufolini MG, Pippi L, Corbisiero R, Fiorentini C, Gistri A, Toscano L. Phlebotomus-transmitted toscana virus infections of the central nervous system: a seven-year experience in Tuscany. *Scand J Infect Dis* 1998; **30**(5):505–8.
- Papa A, Kontana A, Tsergouli K. Phlebovirus infections in Greece. *J Med Virol* Jul. 2015; **87**(7):1072–6.
- Mendoza-Montero J, Gamez-Rueda M-I, Navarro-Mari J-M, de la Rosa-Fraile M, Oyonarte-Gomez S. Infections due to sand fly fever virus serotype Toscana in Spain. *Clin Infect Dis Sep.* 1998; **27**(3):434–6.
- Braito A, Corbisiero R, Corradini S, Fiorentini C, Ciufolini MG. Toscana virus infections of the central nervous system in children: a report of 14 cases. *J Pediatr Jan.* 1998; **132**(1):144–8.
- Santos L, Simões J, Costa R, Martins S, Lecour H. Toscana virus meningitis in Portugal, 2002–2005. *Eur Surveill Jun.* 2007; **12**(6):E3–4.
- Anagnostou V, Pardalos G, Athanasiou-Metaxa M, Papa A. Novel phlebovirus in febrile child, Greece. *Emerg Infect Dis May* 2011; **17**(5):940–1.
- Punda-Polic V, Mohar B, Duh D, Bradarić N, Korva M, Fajš L, et al. Evidence of an autochthonous Toscana virus strain in Croatia. *J Clin Virol Sep.* 2012; **55**(1):4–7.
- Dupouey J, Bichaud L, Ninove L, Zandotti C, Thirion-Perrier L, de Lamballerie X, Charrel RN. Toscana virus infections: a case series from France. *J Infect Mar.* 2014; **68**(3):290–5.
- Mellace F, et al. Meningiti, meningo-encefaliti ed encefaliti da virus Toscana in Italia, 2016–2021. Punta dell'iceberg di una arbovirosi endemica poco conosciuta. *Boll Epidemiol Naz* 2023; **3**(2):10–9.
- Jaijakul S, Arias CA, Hossain M, Arduino RC, Wootton SH, Hasbun R. Toscana meningoencephalitis: a comparison to other viral central nervous system infections. *J Clin Virol Nov.* 2012; **55**(3):204–8.
- Baldelli F, Ciufolini MG, Francisci D, Marchi A, Venturi G, Fiorentini C, et al. Unusual presentation of life-threatening Toscana virus meningoencephalitis. *Clin Infect Dis Feb.* 2004; **38**(4):515–20.
- Oechtering J, Petzold GC. Acute hydrocephalus due to impaired CSF resorption in Toscana virus meningoencephalitis. *Neurology Aug.* 2012; **79**(8):829–31.
- Calisher CH, Weinberg AN, Muth DJ, Lazicki JS. Toscana virus infection in United States citizen returning from Italy. *Lancet Jan.* 1987; **1**(8525):165–6.
- Epelboin L, Hausfater P, Schuffenecker I, Riou B, Zeller H, Bricaire F, Bossi P. Meningoencephalitis due to Toscana virus in a French traveler returning from Central Italy. *J Travel Med Sep.* 2008; **15**(5):361–3.
- Sanbonmatsu-Gómez S, Pérez-Ruiz M, Collao X, Sánchez-Seco MP, Morillas-Márquez F, de la Rosa-Fraile M, et al. Toscana virus in Spain. *Emerg Infect Dis Nov.* 2005; **11**(11):1701–7.
- Kay MK, Gibney KB, Riedo FX, Kosoy OL, Lanciotti RS, Lambert AJ. Toscana virus infection in American traveler returning from Sicily, 2009. *Emerg Infect Dis Sep.* 2010; **16**(9):1498–500.
- Greco F, Mauro MV, Tenuta R, Apuzzo G, Giraldo C. A new case of meningitis due to Toscana virus; 2012.
- Suardi LR, Di Lauria N, Pozzi M, Rogasi PG, Barilaro A, Azzolini F, et al. Acute cerebellar ataxia: a rare Toscana virus (TOSV) meningoencephalitis complication. *Int J Neurosci Mar.* 2019; **130**(3):276–8. <https://doi.org/10.1080/00207454.2019.1673748>
- Matusali G, D'Abramo A, Terrosi C, Carletti F, Colavita F, Vairo F, et al. Infectious Toscana virus in seminal fluid of young man returning from Elba Island, Italy. *Emerg Infect Dis* 2022 Apr; **28**(4):865–9.
- Howell BA, Azar MM, Landry ML, Shaw AC. Toscana virus encephalitis in a traveler returning to the United States. *J Clin Microbiol Apr.* 2015; **53**(4):1445–7.
- Ocal M, Orsten S, Inkaya AC, Yetim E, Acar NP, Alp S, et al. Ongoing activity of Toscana virus genotype A and West Nile Virus lineage 1 strains in Turkey: a clinical and field survey. *Zoonoses Public Health Nov.* 2014; **61**(7):480–91.
- Bartels S, Boni L, Kretzschmar HA, Heckmann JG. Lethal encephalitis caused by the Toscana virus in an elderly patient. *J Neurol Jan.* 2012; **259**(1):175–7.
- Martínez-García FA, Moreno-Docón A, Segovia-Hernández M, Fernández-Barreiro A. Deafness as a sequela of Toscana virus meningitis. *Med Clin May* 2008; **130**(16):639.
- Osborne JC, Khatamzas E, Misbahuddin A, Hart R, Sivaramakrishnan A, Breen DP. Toscana virus encephalitis following a holiday in Sicily. *Pract Neurol Apr.* 2016; **16**(2):139–41.
- Serata D, Rapinesi C, Del Casale A, Simonetti A, Mazzarini L, Ambrosi E, et al. Personality changes after Toscana virus (TOSV) encephalitis in a 49-year-old man: a case report. *Int J Neurosci Jan.* 2011; **121**(3):165–9.
- Ocal M, Orsten S, Inkaya AC, Yetim E, Acar NP, Alp S, et al. Ongoing activity of Toscana virus genotype A and West Nile Virus lineage 1 strains in Turkey: a clinical and field survey. *Zoonoses Public Health Nov.* 2014; **61**(7):480–91.
- Rota E, Morelli N, Immovilli P, De Mitri P, Guidetti D. Guillain-Barré-like axonal polyneuropathy associated with Toscana virus infection: a case report. *Medicine* 2017 Sep; **96**(38):8081.

40. Okar SV, Bekircan-Kurt CE, Hacıoğlu S, Erdem-Özdamar S, Özkul A, Ergünay K. Toscana virus associated with Guillain-Barré syndrome: a case-control study. *Acta Neurol Belg* Jan. 2020;**121**(3):661–8.
41. Tappe D, Schmidt-Chanasit J, Günther S, Ries A, Ziegler U, Müller A, Stich A. Acute Toscana virus infection mimicked by Yersinia-induced reactive arthritis syndrome after journey to Spain. *J Clin Virol* Jan. 2010;**47**(1):104–5.
42. Echevarría J-M, de Ory F, Guisasaola ME, Sánchez-Seco MP, Tenorio A, Lozano A, et al. Acute meningitis due to Toscana virus infection among patients from both the Spanish Mediterranean region and the region of Madrid. *J Clin Virol* Jan. 2003;**26**(1):79–84.
43. Anagnostou V, Pardalos G, Athanasiou-Metaxa M, Papa A. Novel phlebovirus in febrile child, Greece. *Emerg Infect Dis* May 2011;**17**(5):940–1.
44. Zanelli G, Bianco C, Cusi MG. Testicular involvement during Toscana virus infection: an unusual manifestation? *Infection* Jun. 2013;**41**(3):735–6.
45. Schultze D, Korte W, Rafeiner P, Niedrig M. First report of sand fly fever virus infection imported from Malta into Switzerland, October 2011. *Eur Surveill* 2012;**17**(27):20209.
46. Arden KE, Heney C, Shaban B, Nimmo GR, Nissen MD, Sloots TP, Mackay IM. Detection of Toscana virus from an adult traveler returning to Australia with encephalitis. *J Med Virol* Oct. 2017;**89**(10):1861–4.
47. Dominati A, Sap L, Vora S. Fever in a returning traveler from Tuscany. *Rev Med Suisse* Jan. 2018;**14**(592):294–6.
48. Vilibic-Cavlek T, Zidovec-Lepej S, Ledina D, Knezevic S, Savic V, Tabain I, et al. Clinical, virological, and immunological findings in patients with Toscana neuroinvasive disease in Croatia: report of three cases. *Trop Med Infect Dis* Sep. 2020;**5**(3):144.
49. Gámbaro F, Pérez AB, Prot M, Agüera E, Baidaliuk A, Sánchez-Seco MP, et al. Untargeted metagenomic sequencing identifies Toscana virus in patients with idiopathic meningitis, southern Spain, 2015 to 2019. *Eur Surveill* 2023 Nov;**28**(45):2200913.
50. Erdem H, Ergunay K, Yilmaz A, Naz H, Akata F, Inan AS, et al. Emergence and co-infections of West Nile virus and Toscana virus in Eastern Thrace, Turkey. *Clin Microbiol Infect* Apr. 2014;**20**(4):319–25.
51. Schirmer L, Wölfel S, Georgi E, Ploner M, Bauer B, Hemmer B. Extensive recruitment of plasma blasts to the cerebrospinal fluid in Toscana virus encephalitis. *Open Forum Infect Dis* 2015 Sep;**2**(3):124.
52. Popescu CP, Cotar AI, Dinu S, Zaharia M, Tardei G, Ceausu E, et al. Emergence of Toscana virus, Romania, 2017–2018. *Emerg Infect Dis* 2021 May;**27**(5):1482–5.
53. Quattrone F, Mazzetti P, Aquino F, Sani S, Carneglia L, Pistello M, et al. Two clusters of Toscana virus meningo-encephalitis in Livorno Province and Elba Island, July–September 2018. *Ann Ig Nov*. 2020;**32**(6):674–81.
54. Nicoletti L, Verani P, Cacioli S, Ciufolini MG, Renzi A, Bartolozzi D, et al. Central nervous system involvement during infection by phlebovirus Toscana of residents in natural foci in Central Italy (1977–1988). *Am J Trop Med Hyg* Oct. 1991;**45**(4):429–34.
55. Pauli C, Schwarz TF, Meyer CG, Jager G. Neurological symptoms after an infection by the sand fly fever virus. *Dtsch Med Wochenschr* 1995;**120**(43):1468–72.
56. Mosnier E, Charrel R, Vidal B, Ninove L, Schleinitz N, Harlé JR, Bernit E. Toscana virus myositis and fasciitis. *Med Mal Infect* May 2013;**43**(5):208–10.
57. Tschumi F, Schmutz S, Kufner V, Heider M, Pigny F, Schreiner B, et al. Meningitis and epididymitis caused by Toscana virus infection imported to Switzerland diagnosed by metagenomic sequencing: a case report. *BMC Infect Dis* Jul. 2019;**19**(1):591.
58. Mascitti H, Calin R, Dinh A, Makhoulfi S, Davidov B. Testicular pain associated with clear fluid meningitis: how many cases of Toscana virus are we missing? *Int J Infect Dis* Apr. 2020;**93**:198–200.
59. Kuhn J, Bewermeyer H, Hartmann-Klosterkoetter U, Emmerich P, Schilling S, Valassina M. Toscana virus causing severe meningoencephalitis in an elderly traveller. *J Neurol Neurosurg Psychiatry* Nov. 2005;**76**(11):1605–6.
60. Braito A, Corbisiero R, Corradini S, Marchi B, Sancasciani N, Fiorentini C, Ciufolini MG. Evidence of Toscana virus infections without central nervous system involvement: a serological study. *Eur J Epidemiol* 1997;**13**(7):761–4.
61. Dionisio D, Valassina M, Ciufolini MG, Vivarelli A, Esperti F, Cusi MG, et al. Encephalitis without meningitis due to sand fly fever virus serotype Toscana. *Clin Infect Dis* Apr. 2001;**32**(8):1241–3.
62. Gabriel M, Resch C, Günther S, Schmidt-Chanasit J. Toscana virus infection imported from Elba into Switzerland. *Emerg Infect Dis* Jun. 2010;**16**(6):1034–6.
63. Serata D, Rapinesi C, Del Casale A, Simonetti A, Mazzarini L, Ambrosi E, et al. Personality changes after Toscana virus (TOSV) encephalitis in a 49-year-old man: a case report. *Int J Neurosci* Jan. 2011;**121**(3):165–9.
64. Di Nicuolo G, Pagliano P, Battisti S, Starace M, Mininni V, Attanasio V, Faella FS. Toscana virus central nervous system infections in Southern Italy. *J Clin Microbiol* 2005;**43**(12):6186–8.
65. Doudier B, Ninove L, Million M, de Lamballerie X, Charrel RN, Brouqui P. Unusual Toscana virus encephalitis in southern France. *Med Mal Infect* Jan. 2011;**41**(1):50–1.
66. Marlinge M, et al. A febrile meningoencephalitis with transient central facial paralysis due to Toscana virus infection, south-eastern France. *Eur Surveill* 2014;**19**(48):20974.
67. Sanbonmatsu-Gámez S, Pérez-Ruiz M, Palop-Borrás B, Navarro-Marí JM. Unusual manifestation of toscana virus infection, Spain. *Emerg Infect Dis* Feb. 2009;**15**(2):347–8.
68. Schwarz TF, Jager G, Gilch S, Pauli C. Serosurvey and laboratory diagnosis of imported sand fly fever virus, serotype Toscana, infection in Germany. *Epidemiol Infect* 1995;**114**:501–10.
69. Laroche L, Jourdain F, Ayhan N, Bañuls AL, Charrel R, Prudhomme J. Incubation period for neuroinvasive Toscana virus infections. *Emerg Infect Dis* 2021 Dec;**27**(12):3147–50.
70. Masse S, Ayhan N, Capai L, Bosseur F, de Lamballerie X, Charrel R, Falchi A. Circulation of Toscana virus in a sample population of Corsica, France. *Viruses* Sep. 2019;**11**(9):817.
71. McDonald E, Mathis S, Martin SW, Staples J, Erwin, Fischer M, Lindsey NP. Surveillance for West Nile Virus disease – United States, 2009–2018. *MMWR Surveill Summ* 2021;**70**(1):1–15.
72. Worku DA. Tick-borne encephalitis (TBE): from tick to pathology. *J Clin Med* 2023;**12**(21):6859.
73. Oliveira ARS, Cohnstaedt LW, Noronha LE, Mitzel D, McVey DS, Cernicchiaro N. Perspectives regarding the risk of introduction of the Japanese encephalitis virus (JEV) in the United States. *Front Vet Sci* 2020;**7**:48.
74. Hemmersbach-Miller M, Parola P, Charrel RN, Paul Durand J, Brouqui P. Sand fly fever due to Toscana virus: an emerging infection in southern France. *Eur J Intern Med* Aug. 2004;**15**(5):316–7.
75. Kuhn J, Bewermeyer H, Hartmann-Klosterkoetter U, Emmerich P, Schilling S, Valassina M. Toscana virus causing severe meningoencephalitis in an elderly traveller. *J Neurol Neurosurg Psychiatry* Nov. 2005;**76**(11):1605–6.
76. Oechtering J, Petzold GC. Acute hydrocephalus due to impaired CSF resorption in Toscana virus meningoencephalitis. *Neurology* Aug. 2012;**79**(8):829–31.
77. Epelboin L, Hausfater P, Schuffenecker I, Riou B, Zeller H, Bricaire F, Bossi P. Meningoencephalitis due to Toscana virus in a French traveler returning from Central Italy. *J Travel Med* Sep. 2008;**15**(5):361–3.
78. Zanelli G, Bianco C, Cusi MG. Testicular involvement during Toscana virus infection: an unusual manifestation? *Infection* Jun. 2013;**41**(3):735–6.
79. Kvam KA, Stahl JP, Chow FC, Soldatos A, Tattevin P, Sejvar J, Mailles A. Outcome and sequelae of infectious encephalitis. *J Clin Neurol* 2024 Jan;**20**(1):23–36.
80. Francisci D, Papili R, Camanni G, Morosi S, Ferracchiato N, Valente M, et al. Evidence of Toscana virus circulation in Umbria: first report. *Eur J Epidemiol* 2003;**18**(5):457–9.
81. Vocale C, Bartoletti M, Rossini G, Macini P, Pascucci MG, Mori F, et al. Toscana virus infections in Northern Italy: laboratory and clinical evaluation. *Vector Borne Zoonotic Dis* Jun. 2012;**12**(6):526–9.